

Amendments to the Claims:

The listing of claims will replace all prior versions and listings of claims in the application:

Listing of Claims:

Claim 1 (Original) Process for the preparation of a water continuous acidified emulsion comprising 15 to 50 wt% of a partly crystalline fat and 1 to 6 wt% protein, wherein the process comprises the sequence of

- a) homogenising at a temperature below 60°C, a water continuous emulsion comprising a partly crystalline fat and protein wherein the percentage of denatured protein on total protein is less than $100 - (2.6 * a/b)$, wherein a is the amount of fat on the total product in wt% and b is the amount of total protein on the product in wt%,
- b) heating to a temperature above 80°C for a time longer than 5 minutes, and
- c) acidifying to a pH below the pH of gelling of the protein having the highest pH of gelling in the emulsion.

Claim 2 (Original) Process according to claim 1 wherein the emulsion comprises 15 to 25 wt% of a partly crystalline fat and 2 to 3 wt% protein, wherein the process comprises the sequence of

- a) homogenising at a temperature below 60°C, a water continuous emulsion comprising a partly crystalline fat and protein wherein the percentage of denatured protein on total protein is less than 67 %,
- b) heating to a temperature above 85°C for a time longer than 10 minutes, and
- c) acidifying to a pH below the pH of gelling of the protein having the highest pH of gelling in the emulsion.

| Claim 3 (Currently Amended) Process according to claim 1 or 2 wherein the emulsion before homogenisation is prepared by mixing fat and protein, in an aqueous phase.

Claim 4 (Original) Process for the preparation of a water continuous acidified emulsion comprising 15 to 50 wt% partly crystalline fat and 1 to 6 wt% protein, wherein the process comprises the sequence of

- a) homogenising at a temperature below 60°C, a water continuous emulsion comprising fat and protein wherein the percentage of denatured protein is less than 20% on total protein.
- b) heating to a temperature and for sufficient time to prepare an emulsion wherein the percentage of denatured protein is more than 20% on total protein and
- c) acidifying to a pH below the pH of gelling of the protein having the highest pH of gelling in the emulsion.

Claim 5 (Original) Process according to claim 4 wherein the emulsion before homogenisation is prepared by mixing fat and protein, in an aqueous phase where the percentage of denatured protein is less than 20% on total protein.

| Claim 6 (Currently Amended) Process according to ~~any of claim 1 to 5~~ claim 1 wherein the fat is a vegetable fat.

| Claim 7 (Currently Amended) Process according to ~~any of claim 1 to 6~~ claim 1, wherein the partially crystalline fat has a fatty acid composition wherein the combined amount of HM2, M3, MSh2, M2Sh, and HMSh is at least 46 wt% and the amount of U3 is less than 28 wt%, and the combined amount of U3 and H2U is less than 30 wt%, based on total fatty acids.

Claim 8 (Original) Process according to claim 7 wherein the partly crystalline fat comprises least 50 wt% of fully hardened coconut fat or at least 55 wt% of fully hardened palm kernel oil or combinations of x wt% of fully hardened coconut fat and y wt% of fully hardened palm kernel oil, wherein $x + (50/55)y$ is at least 50, based on weight of fat.

| Claim 9 (Currently Amended) Process according to ~~any of claim 1 to 8~~claim 1 wherein
the product of step (b) is acidified in step (c) to a pH in the range of 5.8 to 3.8.

| Claim 10 (Currently Amended) A water continuous acidified emulsion obtainable by the
process according to ~~any of claims 1 to 9~~claim 1.

| Claim 11 (Currently Amended) A process according to ~~any of claims 1 to 9~~claim 1 or a
product according to claim 10 wherein the protein is a whey protein.